


1/17


Figure 1

Adsorption isotherms of benzene on ZSM-5 and Y
Zeolite




2/17

Figure 2



3/17


Figure 3
One column system



4/17

Figure 4


Two column system



5/17


Figure 5

Gaschromatographic profile of the head space (GC-MS)
of groundwater, in the feeding to the PRB simulation
apparatus



6/17


Figure 6
Gaschromatographic profile of the head space (GC-MS)
of groundwater, at the outflow of the PRB simulation
apparatus



7/17


Figure 7

Gaschromatographic profile of the head space (GC-MS)
of water removed from the water-bearing stratum



8/17

Figure 8



9/17


Figure 9

Treatment of real water

Column (7,5x40cm)

0-20 cm Zeolite Y (50g); 21-40 cm ZMS-5 (50g)

Breakthrough at 20 cm (Y Zeolite)



10/17


Figure 10

Treatment of eal water

Column (7.5x40cm)



0-20 cm Y Zeolite (50g); 21-40 cm ZMS-5 (50g)

Breakthrough at 40 cm (ZMS-5)



11/17


Figure 11



13/17


Figure 13

Gaschromatographic profile of the head space of water
at the outflow of the Y zeolite column




14/17

Figure 14




15/17

Figure 15



16/17

Figure 16



17/17

Figure 17

